



XXVIIth International Conference on Ultrarelativistic Nucleus-Nucleus Collisions
(Quark Matter 2018)

Heavy-flavour hadron decay leptons in Pb–Pb and Xe–Xe collisions at the LHC with ALICE

Andrea Dubla for the ALICE Collaboration

*Physikalisches Institut, Universität Heidelberg, 69120 Heidelberg, Germany
and GSI Helmholtzzentrum für Schwerionenforschung, 64248 Darmstadt, Germany*

Abstract

Heavy quarks, i.e. charm and beauty, are sensitive probes of the medium produced in high-energy heavy-ion collisions. They are produced in the early stage of the collision, mainly in hard partonic scattering processes, and are expected to experience the whole collision evolution interacting with the medium constituents via both elastic and inelastic processes. The nuclear modification factor (R_{AA}) is one of the main experimental observables that allow us to investigate the interaction strength of heavy quarks with the medium. The ALICE collaboration measured the production of open heavy-flavour hadrons via their semi-leptonic decays to electrons at mid-rapidity and to muons at forward rapidity in elementary proton-proton (pp) collisions as well as p–Pb, Pb–Pb and in Xe–Xe collisions. Recent results will be discussed, and model calculations including the interaction of heavy quarks with the hot and dense QCD medium will be compared with the data.

Keywords:

ALICE, heavy-flavour, heavy-ion

1. Introduction

The main goal of the ALICE experiment [1] is to study the strongly-interacting matter at high energy density and temperature produced in ultra-relativistic heavy-ion collisions at the Large Hadron Collider (LHC). In such collisions a deconfined state of quarks and gluons, the Quark-Gluon Plasma (QGP), is expected to be formed. Due to their large masses, heavy quarks, i.e. charm (c) and beauty (b) quarks, are produced at the initial stage of the collision, almost exclusively in hard partonic scattering processes. Therefore, witnessing the full evolution of the system, they are effective probes to study the mechanisms of parton energy loss [2, 3] and hadronisation [4] in the hot and dense medium, giving insight on the QGP evolution and its transport coefficients. Finally, initial-state effects due to the presence of a heavy nucleus in the collision system can play a role. At low Bjorken- x (below 10^{-2}) the parton densities in nucleons bound in nuclei are reduced with respect to those in free nucleons. This so-called "shadowing" leads to a reduction of heavy-flavour production rates, becoming more pronounced with decreasing p_T [5]. Initial-state effects and their impact on the nuclear modification factor are investigated in p–Pb collisions. In order to exploit

the sensitivity of heavy-flavour observables to medium effects a precise reference where such effects are not expected is needed and it is provided by pp collisions. In pp collisions, heavy-quark production can be described theoretically via perturbative QCD calculations over the full quark momentum range, while such a description does not hold for gluon and light-quark production. Therefore, measurements of heavy-flavour production cross sections in pp collisions are used to test perturbative QCD calculations and provide the necessary experimental reference for heavy-ion collisions. The modification of the p_T -differential yield in heavy-ion collisions with respect to pp collisions at the same centre-of-mass energy is quantified by the nuclear modification factor R_{AA} , defined as: $R_{AA}(p_T, y) = \frac{1}{\langle T_{AA} \rangle} \cdot \frac{d^2 N_{AA}/dp_T dy}{d^2 \sigma_{pp}/dp_T dy}$, where $d^2 N_{AA}/dp_T dy$ is the yield measured in heavy-ion collisions in a given p_T and y interval, and $d^2 \sigma_{pp}/dp_T dy$ is the corresponding production cross section in pp collisions.

2. Heavy-flavour hadron decay leptons in pp, Xe–Xe and Pb–Pb collisions

The p_T -differential production cross section of electrons from heavy-flavour hadron decays has been measured at mid-rapidity in pp collisions over a broad range of collision energies and it is shown in Fig. 1 (from left to right panel: $\sqrt{s} = 2.76, 5.02, 7$ and 13 TeV). The photonic-tagging background subtraction method [7] allowed for a reduction of the systematic uncertainties on the pp reference cross section by a factor of about 3 compared to the previously published cross sections at $\sqrt{s} = 2.76$ and 7 TeV [8, 9]. The dashed areas represent the FONLL calculation [6]. The full systematic band of the model originates from the variation of the factorization and normalization scales as well as the heavy-quark masses and the uncertainty on the parton distribution function (PDF) used. In the lower panels of Fig. 1 the ratios of the experimental measurements with the central value of the FONLL calculations are shown. The measured cross sections are close to the upper edge of the FONLL uncertainty band for all the collision energies analysed.




Fig. 1. p_T -differential production cross section for electrons from heavy-flavour hadron decay in pp collisions at several collision energies (from left to right panel: $\sqrt{s} = 2.76, 5.02, 7$ and 13 TeV) in comparison with FONLL calculations [6].

The ALICE collaboration measured the R_{AA} of open heavy-flavour hadrons via their semi-leptonic decays in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ [10] and 5.02 TeV. The left panel of Fig. 2 shows the R_{AA} of electrons from heavy-flavour hadron decays as a function of p_T in central (0–10%) Pb–Pb and in p–Pb collisions [7] at $\sqrt{s} = 5.02$ TeV. A strong suppression is observed in central Pb–Pb collisions above $p_T = 1$ GeV/c, while an R_{pPb} consistent with unity is measured, confirming that the suppression observed in central Pb–Pb collisions is predominantly induced by final-state effects due to the heavy quark energy loss in the medium. In the low p_T region, the nuclear modification of the PDFs can play a significant role in the production of heavy quarks. This is addressed in the right panel of Fig. 2, which compares the measured R_{AA} of electrons from heavy-flavour hadron decays at $\sqrt{s} = 2.76$ TeV with TAMU, POWLANG and MC@sHQ+EPOS2 model calculations [11, 12, 13] with and without the inclusion of the EPS09 shadowing

parameterisations [5]. The depletion of the parton densities at low x , resulting in a reduced heavy-flavour production cross section per nucleon-nucleon pair in Pb–Pb collisions with respect to bare nucleon–nucleon collisions, leads to a reduction of the R_{AA} of electrons from heavy-flavour hadron decays at low p_T . Data are better described when the nuclear PDFs are included in the theoretical calculations.




Fig. 2. Left: R_{AA} of electrons from heavy-flavour hadron decays as a function of p_T in central (0–10%) Pb–Pb (black and grey markers) and in p–Pb (blue markers) collisions [7] at $\sqrt{s} = 5.02$ TeV. Right: R_{AA} of electrons from heavy-flavour hadron decays at $\sqrt{s} = 2.76$ TeV in comparison with model calculations with and without the inclusion of the EPS09 shadowing parameterisations [11, 12, 13].

Due to the QCD nature of parton energy loss, quarks are predicted to lose less energy than gluons due to color-charge effect (smaller coupling with the plasma). In addition, the dead-cone effect [14] is expected to reduce the energy loss of massive quarks with respect to light quarks. Therefore, if we consider only energy-loss effects, a hierarchy in the R_{AA} is expected to be observed when comparing the mostly gluon-originated light-flavour hadrons (e.g. pions) to D and to B mesons. The measurement and comparison of these different medium probes provide a unique test of the colour-charge and mass dependence of parton energy loss. A comparison of the R_{AA} of electrons from beauty-hadron decays with the one from charm-plus beauty-hadron decays is shown in the left panel of Fig. 3 for the 10% most central Pb–Pb collisions at $\sqrt{s} = 5.02$ TeV. The contribution to the heavy-flavour decay electron yield due to beauty-hadron decays was extracted by means of a fit to the electron impact parameter distribution. The two R_{AA} are fully compatible for $p_T > 6$ GeV/c, where the beauty contribution is larger than the charm one [8]. At lower p_T , where the charm contribution dominates, the data indicates a smaller suppression for electrons coming from beauty-hadron decays with respect to electrons from inclusive heavy-flavour hadron decays. The measured R_{AA} is in agreement within uncertainties with models implementing mass-dependent energy loss [12, 15, 16].

The nuclear modification factor of electrons (and muons) from heavy-flavour hadron decays in Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV in the kinematic range $0.2(3) < p_T < 6(8)$ GeV/c are reported for the centrality classes 0–20% (0–10%) and 20–40% in the left and right panel of Fig. 4, respectively. A pp reference at $\sqrt{s} = 5.44$ TeV is obtained by the extrapolation of the existing spectra at $\sqrt{s} = 5.02$ TeV with a FONLL based scaling. The R_{AA} values of electron and muons from heavy-flavour hadron decays are comparable in magnitude. When comparing nuclear modification factors at similar ranges of averaged charged particle multiplicity densities, a remarkable similarity between central Xe–Xe collisions and Pb–Pb collisions at a similar center-of-mass energy is observed. The comparison of the measured R_{AA} values in the two colliding systems could enable a test of the path length dependence of medium-induced parton energy loss.

3. Conclusion

Measurements of production of open heavy-flavour hadrons via their semi-leptonic decays to electrons at mid-rapidity and to muons at forward rapidity have been presented in pp, p–Pb, Pb–Pb and Xe–Xe collisions. The results in Pb–Pb and in Xe–Xe collisions indicate a strong suppression of heavy-flavour decay leptons and of electrons from beauty-hadron decays with respect to elementary pp collisions. From the comparison




Fig. 3. Left: Comparison of the R_{AA} of electrons from beauty-hadron decays (red markers) with the one from charm- plus beauty-hadron decays (black markers) in Pb-Pb collisions at $\sqrt{s} = 5.02$ TeV. Right: R_{AA} of electrons from beauty hadron decay in comparison with models implementing mass-dependent energy loss [12, 15, 16] in Pb-Pb collisions at $\sqrt{s} = 5.02$ TeV.




Fig. 4. Nuclear modification factors of electrons (black markers) and muons (red markers) from heavy-flavour hadron decay in central (left panel) and semi-central (right panel) Xe-Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV.

with p-Pb measurements it is possible to conclude that the suppression observed in Pb-Pb and Xe-Xe collisions is mainly due to final state effects.

References

- [1] K. Aamodt *et al.* [ALICE Collaboration], Journal of Instrumentation 3 no. 08, (2008) S08002
- [2] M. Gyulassy and M. Plumer, Phys. Lett. B 243 (1990), 432 - 438
- [3] M.Thoma and M. Gyulassy, Nuclear Physics B 351 (1991), 491 - 506
- [4] V. Greco *et al.* , Phys. Lett. B595 (2004) 202-208
- [5] K. J. Eskola *et al.* , JHEP 04 (2009) 065
- [6] M. Cacciari *et al.* , JHEP 05 (1998) 007
- [7] J. Adam *et al.* [ALICE Collaboration], Phys. Lett. B 754 (2016) 81-93
- [8] B. Abelev *et al.* [ALICE Collaboration], Phys. Rev. D 91 (2015) 226 012001
- [9] B. Abelev *et al.* [ALICE Collaboration], Phys. Rev. D 86 (2012) 112007
- [10] A. Shreyasi *et al.* [ALICE Collaboration], arXiv:1805.04379 [nucl-ex]
- [11] Andrea Beraudo *et al.* , JHEP 02 (2018) 043
- [12] M. Nahrgang *et al.* , Phys. Rev. C 89 (2014), 014905
- [13] M. He *et al.* , Phys. Lett. B 735 (2014), 445 - 450
- [14] Y. L. Dokshitzer and D. E. Kharzeev, Phys. Lett. B 519 (2001), 199 - 206
- [15] M. Djordjevic, Phys. Rev. Lett. B 737 (2014) 286 - 298
- [16] Taesoo Song *et al.* , Phys. Rev. C 93, 034906 (2016)